


# EmergencyKT: Meningitis



## Box 2<sup>5</sup>: CSF Interpretation

Typical Values in various CNS infections\*

Index	Normal	Bacterial	Viral	Fungal	TB
WBC/mcl	<5	>1,000 (100-10,000)	5-1000	<1000	5-1000
Differential	<15% PMNs	>80% PMNs	<15% PMNs (>50% lymphs)	>50% lymphs	>80% lymphs
Glucose (mg/dL)	45-65	reduced	normal	reduced	<40
CSF:Blood glucose	0.6	<0.4	normal	reduced	reduced
Protein (mg/dL)	20-45	>250 (100-500)	30-150	40-150	>50
Opening Pressure (cm/H2O)	<20	elevated	normal or elevated	elevated	variable

Erdman SM, Atkinson KM, Rodvold KA. Chapter 16. Infectious diseases. In: Mary Lee, ed. *Basic Skills in Interpreting Laboratory Data*. 4th ed. Bethesda, MD: American Society of Health-Systems Pharmacists; 2009:391-448.  
 Bahal I, Ellings J. Chronic meningitis. In: Mandell GL, Bennett JE, Dolin R, eds. *Principles and Practice of Infectious Diseases*. 6th ed. Philadelphia, PA: Elsevier Churchill Livingstone; 2005:1132-1143.  
 UpToDate. Cerebrospinal fluid: Physiology and utility of an examination in disease states. Table CSF analysis in CNS infections.

\*significant overlap in findings can occur between various etiologies of meningitis

## Box 3<sup>6</sup>: Steroids

Steroids: to be given prior to or in concomitant with first antibiotic

**Dexamethasone: 0.15 mg/kg  
or 10mg q6hr**

## Box 4<sup>7</sup>: Antibiotics

Neonates (birth-1m)	Children (1m-2y)	Healthy Adults (2y-50y)	Elderly (>50yo)	Immunocompromised	Nosocomial Infections	Healthy Adults – Beta- Lactam Allergy
Ampicillin 50mg/kg q8hr + Cefotaxime 50mg/kg q12hr Or Gentamycin 5mg/kg qday	Vancomycin 15mg/kg q12 hr + Cefotaxime 50mg/kg q12 hr	Vancomycin 15-20 mg/kg q8-12hrs + Ceftriaxone 2g q12-24hr or Cefotaxime 2g q6hr	Vancomycin 15-20mg/kg q8-12hr + Ampicillin 2g q4hr + Cefepime 2g q8hr or Meropenem 2g q8hr	Vancomycin 15-20mg/kg q8-12hr + Ampicillin 2g q4hr + Cefepime 2g q8hr or Meropenem 2g q8hr	Vancomycin 15-20mg/kg q8-12hr + Cefepime 2g q8hr or Ceftazidime 2g q8hr or Meropenem 2g q8hr	Vancomycin 15- 20mg/kg q8-12hr + Meropenem 2g q8hr
Acydovir 20mg/kg q8hr	Acydovir 20mg/kg q8hr (0-3m)  10-5mg/kg (>3m-12y)	Acydovir 10mgkg q8hr (ideal body weight)	Acylovir 10mgkg q8hr (ideal body weight)	Acydovir 10mgkg q8hr (ideal body weight)		Acydovir 10mgkg q8hr (ideal body weight)

Microorganism	Recommended therapy	Alternative therapies
S. pneumonia	Vancomycin+3 <sup>rd</sup> generation cephalosporin	Meropenem, fluoroquinolone
N. meningitidis	3 <sup>rd</sup> generation cephalosporin	Penicillin G, ampicillin, chloramphenicol, aztreonam, fluroquinolone
L. monocytogenes	Ampicillin, penicillin G	Bactrim, meropenem
S. Agalactiae	Ampicillin, penicillin G	3 <sup>rd</sup> generation cephalosporin
H. influenza	3 <sup>rd</sup> generation cephalosporin	Chloramphenicol, cefepime, meropenem, fluoroquinolone
E. Coli	3 <sup>rd</sup> generation cephalosporin	Cefepime, meropenem, aztreonam, Bactrim, fluoroquinolone

## References

1. The rational clinical examination. Does this adult patient have acute meningitis? Attia J, Hatala R, Cook DJ, Wong JG. *JAMA*. 1999;282(2):175.
2. Clinical features and prognostic factors in adults with bacterial meningitis. van de Beek D, de Gans J, Spanjaard L, Weisfelt M, Reitsma JB, Vermeulen M. *N Engl J Med*. 2004;351(18):1849.
3. Hasbun R, Abrahams J, Jekel J, Quagliarello VJ. Computed tomography of the head before lumbar puncture in adults with suspected meningitis. *N Engl J Med* **2001**; 345:1727–33.
4. Sormunen P, Kallio MJT, Kilpi T, Peltola H. C-reactive protein is useful in distinguishing Gram stain–negative bacterial meningitis from viral meningitis in children. *J Pediatr* **1999**; 134:725–9.
5. Durand M.L., Calderwood S.B., Weber D.J., et al. Acute bacterial meningitis in adults: A review of 493 episodes. *N. Engl. J. Med.* 1993;328:21-8.
6. Brouwer MC, McIntyre P, Prasad K, van de Beek. Cochrane Collaboration: Corticosteroids for bacterial meningitis. 2013
7. Tunkel A.R., Hartman B.J., Kaplan S.L., Et al. Infectious Disease society of America practice guidelines for management of bacterial meningitis. *Clin. Infect. Dis.* 2004;39:1264-84