


# EmergencyKT: Cyanide Toxicity



## <sup>1</sup> Cyanide Sources:

- Acetonitrile
- Acrylonitrile
- Cassava Root
- Ground prunus species seeds
- Industrial exposure to cyanide salts
- Electroplating solvents

Suspect Cyanide toxicity in any sudden collapse in lab or industrial settings with cardiovascular instability

<sup>2</sup>**Hydroxocobalamin** will turn skin and secretions red and may affect spectrophotometric testing and LFT's and UA may be deranged after institution of therapy

Pulse oximetry/CO-oximetry may also be affected

## References:

- Baud FJ et al. Value of lactic acidosis in the assessment of the severity of acute cyanide poisoning. Crit Care Med. 2002; 30:2044-2050.
- Borron S et al. Efficacy of Hydroxocobalamin for the Treatment of Acute Cyanide Poisoning in Adult Beagle Dogs. Clin Toxicol. 2006; 44:5-15.
- Borron S et al. Prospective study of Hydroxocobalamin for Acute Cyanide Poisoning in Smoke Inhalation. Ann Emerg Med; 49: 794-801.
- Fortin J-L et al. Prehospital Administration of Hydroxocobalamin for Smoke Inhalation-Associated Cyanide Poisoning: 8 years of Experience in Paris Fire Brigade. Clin Toxicol. 2006; 44:37-44.
- Hall AH et al. Sodium Thiosulfate or Hydroxocobalamin for the Empiric Treatment of Cyanide Poisoning? Ann Emerg Med. 2007; 49: 806-813.
- Uhl W et al. Safety of hydroxocobalamin in healthy volunteers in a randomized, placebo-controlled study. Clin Toxicol. 2006; 44:17-28.